

Total No. of printed pages = 10

3 (Sem 1) CHM M2

2015

**CHEMISTRY**

**(Major)**

Paper : 1.2

Full Marks – 60


Time – Three hours

The figures in the margin indicate full marks for the questions.

1. Answer the following questions (any *seven*) :

1×7=7


(a) Write the IUPAC name of the following compound :




(b) Draw the structural formula of bicycle [4-2-0] octan -3-01.

[Turn over


- (c) Arrange the following species in increasing order of basic strength :



- (d) Suggest an explanation for the difference in dipole moments for the following pair



$$\mu = 2.73 \text{ D}$$



$$\mu = 3.67 \text{ D}$$



- (e) The C-N bond length in  $\text{H}_2\text{NCONH}_2$  is 0.137 nm instead of normal C-N bond length (0.147 nm). Explain.

- (f) Draw and name the tautomeric forms of  $\text{RCH}_2\text{-NO}_2$ .

- (g) Arrange the following compounds in increasing order of nucleophilicity.

Ammonia, pyridine, aniline, potassium amide

- (h) Find whether the following molecules are aromatic, antiaromatic or non-aromatic.



(ii) What is atropisomerism ? Give an example.  $1+1\frac{1}{2}=2\frac{1}{2}$


(d) (i) What do you mean by optical purity ? Illustrate. 2

(ii) Calculate the enantiomeric excess and specific rotation of a mixture containing 10g of (+) -2-butanol and 6g of (-) -2-butanol. The specific rotation of pure (+) -2-butanol is  $+13.5^\circ$ .

4. Answer the following questions :

[Either (i) or (ii) and (iii) or (iv) from (A), (B) and (C).  $10 \times 3 = 30$


(A) (i) Assign R, S-notation to the following chiral molecule : 2



Give an example each of enantiotopic atoms and diastereotopic faces.  $2 \times 1\frac{1}{2} = 3$

Or

- (ii) Following allene is chiral although it has no chiral centre. Explain.



Draw the more stable cis-orientation of cis-1, 3-cyclohexanediol. Which conformation of cis-1, 3-cyclohexanediol is more stable? Explain with due reason.

1+2=3

- (iii) What do you mean by kinetically controlled and thermodynamically controlled reaction? Draw the energy profile diagram for these two reactions.



2+3=5

Or

- (iv) How carbenes are prepared? Between singlet and triplet carbene which one is more stable and why?

1+2=3

Write the product of the following reactions:



20A/3 (Sem 1) CHM M2 (6)

(P)


- (C) (i) Write the product and find a probable mechanism of the following reaction :




Give an example of nucleophilic addition reaction. 2

Or

- (ii) What are Ei elimination reactions ? Discuss the mechanism and stereochemistry of the following reaction :  $1\frac{1}{2} + 2\frac{1}{2} = 4$




Predict the product of the following reaction. 1




Or

(iv) (R)-2-Butanol is found to have lost optical activity after standing in acidic solution. Account for this observation.

Find the product of the following reaction :





Which one of the following will be most reactive in S<sub>N</sub><sup>2</sup> reaction ?



- (B) (i) What product will you obtain when trans-but-2-ene is treated with  $O_3$ ,  $O_4$  and the product is hydrolysed? Find the stereochemistry of the reaction. 5

Or

- (ii) Find the major product of the following reactions:  $2 \times 1\frac{1}{2} = 3$



Which one of the following alkenes will give optically active product with  $Br_2/CCl_4$

- (a) 1-butene (b) propene (c) cis-2-butene (d) trans-2-butene. 2

- (iii) Give a reaction to show evidence that  $SN^1$  mechanism involves carbocation as intermediate.  $2\frac{1}{2}$

How does solvent polarity influence the rate of  $SN^1$  and  $SN^2$  reactions?


$$1\frac{1}{2} + 1 = 2\frac{1}{2}$$

3. Answer the following questions (any three) :  
5×3=15

(a) (i) Between Furan and thiophene which one has higher resonance energy and why?

(ii) Explain why alkynes are less reactive than alkenes towards electrophilic addition reactions.


(b) (i) Which of the following molecules will undergo faster nucleophilic substitution reaction? Explain with reason.



and



(ii) What is neighbouring group participation? Following molecule gives 100% retention of configuration. Explain.




(c) (i) Although in ethyl methyl amine N is asymmetric still it is optically inactive. Explain.



2. Answer the following questions (any four) :  
2×4=8

(a) What is the nature of  $\text{CH}_3\text{CN}$  ? Illustrate with a reaction. 1+1=2

(b) Show the tautomeric forms of the following compound : 1+1=2



Give an example of valence tautomerism.

(c) What are pseudoaromatic compounds ? Give an example. 1+1=2


(d) Arrange the following intermolecular forces in increasing order of strength Dipole-dipole interaction, H-bonding force, London forces. HCl boils at a much higher temperature than Argon. Which forces acting on it ? Explain. 1+1=2

(e) Draw the following molecule :

(i) (E)-2-(But-2-enyl) benzene-1, 3-dicarboxylic acid

(ii) 9-Oxa-6-aza spiro [4, 5] decane 1+1=2

- (iii) What is Cope reaction ? Find the product of the following reaction showing favourable conformation for elimination in Newman projection.



What is E1cb reaction ?

Or

- (iv) Write the different factors on which E1 reactions complete with SN<sup>1</sup> reaction. Illustrate your answer with examples.