

2013

PHYSICS

(Major)

Paper : 3.1

Full Marks : 60

Time : 2½ hours

The figures in the margin indicate full marks
for the questions

GROUP—A

(**Mathematical Methods**)

(Marks : 25).

1. Answer the following : 1×3=3

(a) Define self-adjoint matrix.

(b) Define trace of a matrix.

(c) If A is a Hermitian matrix, show that e^{iA} is unitary.

2. Verify that $(AB)^T = B^T A^T$ where

$$A = \begin{bmatrix} 1 & 2 & 3 \\ 3 & -2 & 1 \end{bmatrix} \text{ and } B = \begin{bmatrix} 1 & 2 \\ 2 & 0 \\ -1 & 1 \end{bmatrix} \quad 2$$

3. Answer any two questions out of (a), (b) and (c) :

(a) (i) If A and B are Hermitian matrices, show that $i(AB - BA)$ is also Hermitian. 1

(ii) If

$$A(\theta) = \begin{bmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{bmatrix}$$

then show that

$$A(\theta)A(\phi) = A(\phi)A(\theta) = A(\theta + \phi) \quad 2$$

(iii) Find the value of λ for which the matrix

$$A = \begin{pmatrix} \cos \psi & -\sin \psi & 0 \\ \sin \psi & \cos \psi & 0 \\ 0 & 0 & \lambda \end{pmatrix}$$

will be orthogonal. 2

(b) (i) Show that inverse of the transpose of a square matrix is the transpose of the inverse. 2

(ii) Prove that any two eigenvectors corresponding to two distinct eigenvalues of a unitary matrix are orthogonal. 3

- (c) Show that for rotation of one frame with respect to another frame with uniform angular velocity $\vec{\omega}$, the equation of motion of the particle in rotating coordinate system is given by

$$m \frac{d'^2 \vec{r}}{dt^2} = \vec{F} - 2m(\vec{\omega} \times \frac{d' \vec{r}}{dt}) - m[\vec{\omega} \times (\vec{\omega} \times \vec{r})]$$

Which term represents the Coriolis force term? Write the equation of motion in case of relative rotation between frames with non-uniform angular velocity $\vec{\omega}$.

$$3+1+1=5$$

4. Answer either (a) and (b) or (c) and (d) :

Either

- (a) (i) Prove that the following matrix is unitary :

2

$$\begin{bmatrix} \frac{1}{2}(1+i) & \frac{1}{2}(-1+i) \\ \frac{1}{2}(1+i) & \frac{1}{2}(1-i) \end{bmatrix}$$

- (ii) Find the inverse of the following matrix from the adjoint of it :

3

$$A = \begin{bmatrix} 1 & -1 & 1 \\ 2 & -1 & 0 \\ 1 & 0 & 0 \end{bmatrix}$$

- (b) (i) Show that every square matrix can be uniquely expressed as the sum of a symmetric and a skew-symmetric matrix. 2

- (ii) Solve by matrix method the following system of equations : 3

$$x + y + z = 8$$

$$x - y + 2z = 6$$

$$3x + 5y - 7z = 14$$

Or

- (c) (i) Show that the eigenvalues of diagonal matrix are precisely the elements in the diagonal. 2

- (ii) Given

$$A = \begin{bmatrix} 1 & 1 & 2 \\ 3 & 1 & 1 \\ 2 & 3 & 1 \end{bmatrix}$$

Compute A^{-1} by using the fact that A satisfies its characteristic equation.

- (d) Find the eigenvalues and eigenvectors of the matrix

$$A = \begin{pmatrix} 2 & -2 & 0 \\ -2 & 1 & -2 \\ 0 & -2 & 0 \end{pmatrix}$$

5

(5)

GROUP—B

(**Electrostatics**)

(Marks : 35)

5. Choose the correct option/Answer the following : 1×4=4

(a) When a test charge is brought from infinity along the perpendicular bisector of the dipole, the work done is

- (i) positive
- (ii) zero
- (iii) negative
- (iv) None of the above

(b) For a dipole, electric field varies as

- (i) r^{-2}
- (ii) r^{-3}
- (iii) r^{-1}
- (iv) r^{-4}

(c) The unit of \vec{D} is

- (i) V/m^2
- (ii) coul/m^2
- (iii) V/m
- (iv) coul/m

(d) What is atomic polarisability?

6. Answer the following questions : 2×3=6

(a) Can an electrostatic field have the form $\vec{E} = a(y\vec{a}_x - x\vec{a}_y)$, where a is a constant?

(b) Show that the function

$$\phi = 3x^2 + 8y - 3z^2$$

can represent the electrostatic potential in a charge-free region.

(c) Define relative permittivity. Write down Clausius-Mosotti relation.

7. Find an expression for the electric field at a point on the axis of a uniformly charged disc of radius a and surface charge density σ . Show that the electric field strength at point P inside a spherically symmetric charge distribution is given by

$$E_i = \frac{1}{4\pi\epsilon_0} \frac{qr}{R^3}$$

where R is the radius of the charge distribution and r is the distance of the internal point P from the centre of the charge distribution.

2+3=5

Or

Find an expression for the torque exerted by one dipole on another dipole. Write down the interaction potential energy of two short electric dipoles separated by a distance. If one of the dipoles is inclined at an angle θ_1 to the radius vector joining them, show that in the state of equilibrium, the other dipole would make an angle θ_2 with it given by

$$\tan\theta_2 = -\frac{1}{2}\tan\theta_1 \quad 2+3=5$$

8. Answer any two questions out of (a), (b), (c) and (d) :

(a) (i) By using the concept of electrical multipoles, find an expression for the electrostatic potential due to a volume distribution of charge. 5

(ii) An electron can be assumed to be uniformly charged sphere having a total charge e and radius R_0 . Show that the electrostatic energy of the electron is given by

$$U = \frac{1}{4\pi\epsilon_0} \left(\frac{3}{5} \frac{e^2}{R_0} \right)$$

If this energy is equal to rest energy m_0c^2 of the electron, what must be its radius? 5

- (b) (i) State and prove Poisson's equation in electrostatics. What form does it take when the charge density is zero? By solving Laplace's equation, show that the potential at a distance r from the axis of an infinitely long conducting cylinder of radius a_0 charged with a surface charge density σ is given by

$$\phi = -\frac{a_0\sigma}{\epsilon_0} \ln\left(\frac{r}{a_0}\right)$$

Take the potential of the cylinder to be zero. 2+3=5

- (ii) What is a polar molecule? Show that the electric intensity inside a parallel-plate capacitor whose dielectric constant increases linearly from the value k_1 in one plate to the value k_2 in other plate is given by

$$E(x) = \frac{\sigma / \epsilon_0}{\left[\left(\frac{k_2 - k_1}{d} \right) x + k_1 \right]}$$

where x is the distance from first plate, σ is the surface density of charge on first plate. The dielectric constant of a monatomic gas at NTP is 1.000538. If the gas is placed in an electric field of 30 kV/m, find the induced dipole moment of an atom. 1+2+2=5

(c) (i) What is meant by an electrical image? A point charge is placed outside a grounded conducting sphere. Find an expression for the intensity of the electric field outside the sphere by the method of electrical images. 1+4=5

(ii) Define dielectric susceptibility. A dipole of length $2d$ and strength p is placed at a distance r from the axis of an infinitely long and uniformly charged wire having a charge λ per unit length. If the dipole lies in the plane containing the axis of the wire and the dipole axis is making an angle θ with the radial line from the force on the dipole towards the wire, then prove that

$$\vec{F}_2 - \vec{F}_1 = \frac{p\lambda \cos\theta}{2\pi\epsilon_0 r^2}$$

where distance $r \gg d$. 1+4=5

(d) (i) Deduce the differential form of Gauss' law for dielectrics. Define the term electrical displacement. 4+1=5

(ii) Show that for a point charge $+q$ placed in a homogeneous dielectric medium of infinite extent, the medium reduces the point charge $+q$ to a value q/k , where k is the dielectric constant of the medium. 5
