

Total number of printed pages-8

3 (Sem-6) PHY M 2

2020

PHYSICS

(Major)

Paper : 6·2

Full Marks : 60

Time : Three hours

The figures in the margin indicate full marks for the questions.

(Mathematical Methods-IV)

(Marks : 15)

1. Answer **any two** from the following :

1×2=2

(a) What is the rank of a tensor which represents a quantity that does not change when axes are rotated ?

(b) In an N -dimensional space, how many terms is contained in each expression represented by $A_{ij}^k B_{lr}^q C_{sq}^n$?

(c) Evaluate $\delta_m^i \delta_n^m \delta_i^n$ in 4-dimensional space.

Contd.

2. Answer **any four** from the following:

2×4=8

(a) Show that δ_ν^μ is an invariant tensor and transforms as a mixed tensor of rank two.

(b) If A_{lm}^{jk} is tensor, test and mention type and rank of tensors A_{jk}^{ij} , A_{lm}^{jm} .

(c) Illustrate "The inner product of tensors can be thought of as outer product followed by contraction."

(d) Show that gradient of a scalar field is a covariant vector.

(e) If A_j^i is a mixed tensor of rank two, show that A_i^i is also a tensor.

3. Answer **any one** from the following:

5×1=5

(a) The Cartesian components of the velocity vector of a fluid in motion in a two-dimensional plane are $v_x = x^2$, $v_y = y^2$. Find the polar components of the velocity vector in terms of polar co-ordinates r, θ . 5

(b) The Cartesian components of the acceleration vector are $a_x = \frac{d^2x}{dt^2}$,

$a_y = \frac{d^2y}{dt^2}$, $a_z = \frac{d^2z}{dt^2}$. Find the radial

component a_r of the acceleration vector in spherical polar co-ordinates.

5

(c) (i) Prove that the sum of two tensors of the same type is also a tensor.

3

(ii) If $A_{\lambda\mu}$ is a skew-symmetric tensor, show that

$$(B_\nu^\mu B_\tau^\sigma + B_\tau^\mu B_\nu^\sigma) A_{\mu\sigma} = 0. \quad 2$$

(Solid State Physics)

(Marks : 45)

4. Choose the correct answer from the following:

1×7=7

(a) Crystalline state is a —

(i) low energy state

(ii) high energy state

(iii) medium energy state

(iv) None of the above

(b) Coordination number of NaCl structure is :

(i) 8

(ii) 6

(iii) 10

(iv) 12

(c) In solids the strongest bond is —

(i) ionic

(ii) covalent

(iii) metallic

(iv) hydrogen

(d) According to Quantum theory of free electrons, the molar specific heat of free electron is —

(i) $C_v = \frac{3}{2}Nk$

(ii) $C_v = (0.01)\frac{3}{2}Nk$

(iii) $C_v = (0.01)Nk$

(iv) $C_v = (0.001)\frac{3}{2}Nk$

(e) The magnetic susceptibility χ of a superconductor has —

(i) a positive value

(ii) $\chi \rightarrow 0$ as $T \rightarrow T_c$

(iii) $\chi \rightarrow \infty$ as $T \rightarrow T_c$

(iv) a negative value

(f) Hysteresis is a property of —

(i) paramagnetic substances

(ii) ferromagnetic substances

(iii) diamagnetic substances

(iv) all of them

(g) One Bohr Magneton is equal to —

(i) $9.27 \times 10^{-24} \text{ amp m}^2$

(ii) $9.27 \times 10^{-24} \text{ amp/m}^2$

(iii) $9.27 \times 10^{-24} \text{ amp/cm}^2$

(iv) $9.27 \times 10^{-24} \text{ amp cm}^2$

5. Give very short answers of the following questions : $2 \times 4 = 8$

(a) Calculate the packing factor for SC structure.

(b) Deduce a relation between the density of crystalline material and lattice constant in a cubic lattice.

(c) A paramagnetic material has a magnetic field strength of 10^4 A/m . If the susceptibility of the material at room temperature is 3.7×10^{-3} , calculate the magnetization and flux density of the material.

(d) State Bloch theorem.

6. Give short answers of the following questions : **(any two)**

(a) Write down Bragg's law in X-ray diffraction and define the different terms used in the equation. From the equation estimate the wavelength of X-ray that can be used for analysis of crystal diffraction. What is glancing angle ? $2+2+1=5$

(b) What are Miller indices ? How are they determined ? Explain with the help of an example. $1+4=5$

(c) Explain Meissner effect. Outline some applications of superconductivity. $2+3=5$

(d) What do you mean by p -type and n -type semiconductor ? How does the conductivity of semiconductor vary with temperature ? Show schematically the position of Fermi level at 0K in p -type and n -type semiconductor. $2+1+2=5$

7. Answer the following questions :

(a) What do you mean by cohesive energy ? Evaluate Madelung constant for an infinitely long one-dimensional ionic crystal consisting of singly charged alternate positive and negative ions. State the significance of Madelung constant. $2+6+2=10$

Or

(b) On the basis of Weiss theory, obtain Curie-Weiss law. Show that ferromagnetic substances become paramagnetic above a critical temperature. $8+2=10$

(c) Write short note on: **(any one)** 10

(i) Intrinsic and extrinsic semiconductors

(ii) Kronig-Penney model

(iii) Different types of crystal bonding