

2019

MATHEMATICS

( Major )

Paper : 6.4

( Discrete Mathematics )

Full Marks : 60

Time : 3 hours

*The figures in the margin indicate full marks  
for the questions*

1. Answer the following questions :  $1 \times 7 = 7$

(a) State division algorithm of integers.

(b) Write the least positive integer of the form  $172x + 20y$ ,  $x, y \in \mathbb{Z}$ .

(c) If  $p$  is a prime and  $a \in \mathbb{Z}$ , then show that  $(a, p) = 1$  or  $p|a$ .

( 2 )

(d) State the converse of Fermat's theorem.  
Is it valid?

(e) Write the value of the sum  $\sum_{d|n} \mu(d)$ .

(f) State Chinese remainder theorem.

(g) What is the geometrical interpretation of a Diophantine equation  $f(x, y) = 0$ ?

2. Answer the following questions :  $2 \times 4 = 8$

(a) Show that there is no positive integer  $n$  such that  $0 < n < 1$ .

(b) Show that  $\phi(n)$  is even if  $n > 2$ .

(c) State and prove the converse of Wilson's theorem.

(d) If  $x, y, z$  are primitive, positive, Pythagorean triple, then show that

$$\left( \frac{z-x}{2}, \frac{z+x}{2} \right) = 1$$

where  $x$  is odd.

( 3 )

3. Answer the following questions :  $5 \times 3 = 15$

(a) Let  $a, b \in \mathbb{Z}$ ,  $a$  or  $b \neq 0$ ,  $G = (a, b)$ . Show that  $G = ax_0 + by_0$  for some  $x_0, y_0 \in \mathbb{Z}$ .

Or

Show that every integer  $n > 1$  can be expressed as a product of primes. Find the prime factorization of  $40!$   $3+2=5$

(b) Using Chinese remainder theorem, find the least positive integer which leaves the remainders 1, 6, 2 when divided by 7, 10, 11 respectively.

Or

Let  $n = p_1^{k_1} p_2^{k_2} \dots p_r^{k_r}$  be the prime factorization of a positive integer  $n > 1$ . Show that the positive divisors of  $n$  are precisely the integers of the form  $d = p_1^{a_1} p_2^{a_2} \dots p_r^{a_r}$ , where  $0 \leq a_i \leq k_i$ ,  $i = 1, 2, \dots, r$ .

( 4 )

(c) The linear congruence  $ax \equiv b \pmod{m}$  has a solution if and only if  $(a, m) | b$ . Prove it.

Or

Show that the Diophantine equation  $x^4 + y^4 = z^2$  has no solutions in positive integers.

4. Answer either (a) or (b) : 10

(a) (i) Let  $p$  be a prime and  $\gcd(a, p) = 1$ . Then show that the congruence  $ax \equiv y \pmod{p}$  has a solution  $x_0, y_0$  such that

$$0 < |x_0| < \sqrt{p}, \quad 0 < |y_0| < \sqrt{p} \quad 5$$

(ii) If  $p_n$  is the  $n$ th prime number, then show that the sum

$$\frac{1}{p_1} + \frac{1}{p_2} + \dots + \frac{1}{p_n}$$

is never an integer. 5

( 5 )

(b) (i) Show that Euler's function  $\phi$  is a multiplicative function. 7

(ii) Show that no prime of the form  $4k+3$  is a sum of two squares. 3

5. Answer either (a) or (b) : 10

(a) (i) Give an example of an infinite Boolean algebra. In a Boolean algebra  $B$ , show that

$$a+a=a, \quad a \cdot (a+b) = a, \quad a, b \in B \quad 2+3=5$$

(ii) State and prove the 'principle of duality' in a Boolean algebra. Write down the dual of the proposition  $a+b=0 \Leftrightarrow a=0, b=0$ . 4+1=5

(b) (i) Simplify the Boolean expression

$$(x+y)(x+z)(x'y)'$$

Draw a switching circuit which realizes the Boolean expression  $x+y(z+x'(t+z'))$ . 3+2=5

( 6 )

- (ii) Construct the switching table for the switching function  $f$  represented by the Boolean expression  $xyz + x'(y + z)$ . 5

6. Answer either (a) or (b) : 10

- (a) (i) Define 'logical equivalence'. Prove that if  $\models A$  and  $\models A \rightarrow B$ , then  $\models B$ . 1+4=5

- (ii) Construct the truth tables for NOR( $\downarrow$ ) and NAND( $\uparrow$ ). Show that  $\{\wedge, \rightarrow\}$  is not an adequate system of connectives. 2+3=5

- (b) (i) Using principle of substitution, show that if  $A, B$  be any two statement formulae, then  $A \rightarrow B \equiv \sim B \rightarrow \sim A$  5

- (ii) Assuming the truth value of  $p \rightarrow q$  be  $T$ , construct the truth table for  $(\sim p \wedge q) \leftrightarrow (p \vee q)$ . 2

( 7 )

- (iii) Define a truth function. Construct the truth function generated by the statement formula

$$\sim(\sim p \wedge q) \quad 1+2=3$$

\*\*\*