### Total number of printed pages-7

## xintem llum a 3 (Sem-3/CBCS) PHY HC1

ACC. NO.

2024

#### PHYSICS

(Honours Core)

Paper: PHY-HC-3016

(Mathematical Physics-II)

Full Marks: 60

Time: Three hours

# The figures in the margin indicate full marks for the questions.

- 1. Answer the following questions: 1×7=7
  - (a) Define regular singular point at x = 0 for a second order differential equation.
  - (b) The function  $P_n(1)$  is given as
    - (i) -1
    - (ii) zero
- Using the property of gam (iii) function
  - (iv)  $P_n(-1)$

(Choose the correct option)

- Define gamma function.
- (d) Write the rank of a null matrix.
- (e) What do you mean by unitary matrix?
- Write the orthogonal property of Hermite polynomials.
- State the Dirichlet condition for Fourier series.
- Answer the following questions: The floures in the manuin insteate

for a second order differential equation

- full searcs for the guestions (a) If  $\int_{-1}^{1} P_n(x) dx = 2$ , then find the value Define regular singular in lo at x
- (b) If A and B are Hermitian matrices, show that AB + BA is Hermitian and AB - BA is skew-Hermitian.
- Using the property of gamma function evaluate the integral  $\int_{0}^{\infty} x^{4}e^{-x}dx$ .

- (d) If  $f(x) = x \cos x$  is a function in the interval  $-\pi < x < \pi$ , then find the value of  $a_0$  of the Fourier series.
- Answer any three of the following questions:  $5 \times 3 = 15$ 
  - (a) If the solution y(x) of Hermite's differential equation is written as
    - $y(x) = \sum_{r=0}^{\infty} a_r x^{k+r}$ , then show that the (e) Find the Fourier series repres allowed values of k are zero and one only.
  - (a) (i) Find the eigenvalue Obtain the following orthogonality property of Legendre polynomial

$$\int_{-1}^{+1} P_n(x) P_m(x) dx = 0 \text{ for } m \neq n$$

(c) (i) What is a special square matrix?

interval 
$$-x < \begin{bmatrix} 2 & 2 & 1 \\ 1 & 3 & 1 \end{bmatrix} > x - law value$$

$$A = \begin{bmatrix} 1 & 3 & 1 \\ 1 & 2 & 2 \end{bmatrix} \cdot hod the value$$

$$A = \begin{bmatrix} 1 & 2 & 2 \\ 1 & 2 & 2 \end{bmatrix} \cdot hod the value$$

(d) What is adjoint of a matrix? For the matrix  $A = \begin{bmatrix} 1 & 2 \\ 3 & -5 \end{bmatrix}$  verify the theorem

Answer any three of the following

A.(Adj A) = (Adj A). A = |A|I, where I is unit matrix. 1+4=5

- Find the Fourier series representing one buf f(x) = x,  $0 < x < 2\pi$  they be well a
- (a) (i) Find the eigenvalue and eigenvectors of the matrix

$$A = \begin{bmatrix} b_2 & b_1 \\ -1 & 4 \end{bmatrix}.$$
 6

Show that any arbitrary square matrix can be represented as a sum of symmetric and skewsymmetric matrix.

#### OR

(b) (i) Prove the Rodrigues' formula for Legendre's polynomials

$$P_n(x) = \frac{1}{2^n \cdot n!} \frac{d^n}{dx^n} (x^n - 1)^n$$

Show that wig on lo

$$H_0(x) = 1$$
 and  $H_1(x) = 2x$ .



(a) (i) Solve the following equation by the method of separation of variables

hij Write the orthogonality

$$\frac{\partial u}{\partial x} = 4 \frac{\partial u}{\partial y}, \text{ bins only 10}$$

given 
$$u(0, y) = 8e^{-3y}$$
 5

(ii) Write the recursion formula of gamma function. Prove that

$$\Gamma\left(\frac{1}{2}\right) = \sqrt{\pi}.$$
1+4=5

Show that the matrix (b) (i)

OR

$$A = \begin{bmatrix} \frac{1}{\sqrt{2}} & \frac{i}{\sqrt{2}} \\ -\frac{i}{\sqrt{2}} & -\frac{1}{\sqrt{2}} \end{bmatrix}$$
 is unitary.

3

Find the power series solution of the differential equation

$$\frac{d^2y}{dx^2} + w^2y = 0 \text{ in powers of } x. \quad 7$$



(b) (i) A square wave function is represented as



$$f(x) = \begin{cases} 0, & \text{for } -\pi < x < 0 \\ L, & \text{for } 0 \le x < \pi \end{cases}$$

Draw the graphical representation of the given function. Show that the Fourier expansion of the function is given as

$$f(x) = \frac{L}{2} + \frac{2L}{\pi} \sum_{n=1}^{\infty} \frac{\sin nx}{n}$$
 (for *n* odd)

and you contain by the following equation by the

- (ii) Write the orthogonality conditions of sine and cosine functions.
- 6. (a) Prove the following recurrence relations: 3+2+5=10

method of separation of variables

lo si (i) ol 
$$2xH_n(x) = 2nH_{n-1}(x) + H_{n+1}(x)$$

(ii) 
$$H_n^1(x) = 2x H_n(x) - H_{n+1}(x)$$
  
(iii)  $x P_n^1(x) - P_{n-1}^1(x) = n P_n(x)$ 

(iii) 
$$x P_n^1(x) - P_{n-1}^1(x) = n P_n(x)$$